arXiv:0806.0892v6 [math.CA] 18 Oct 2008

ON ZEROS OF CERTAIN ENTIRE FUNCTIONS

RUIMING ZHANG

Abstract. In this work we prove that a class of entire functions considered by Jensen and Pólya have only real zeros. As a consequence, we may have proved the Riemann Hypothesis for the Riemann zeta function $\zeta(s)$ and the Riemann the character Riemann zeta function $L(\chi, x)$ with real primitive characters satisfying certain positive conditions.

1. Introduction

In this work we study the zeros of certain entire functions. They are of genus at most one, and defined as Fourier transforms of some even nonnegative functions which are fast decreasing on the real line. Jensen and Pólya investigated the properties of these functions in the context of the Riemann Hypothesis. Jensen had proved several necessary and sufficient conditions for the these entire functions having only real zeros [5, 7]. We shall prove that these entire functions have only real zeros by verifying one set of Jensen's necessary and sufficient conditions directly. As a consequence, we may have proved the Riemann Hypotheses for the Riemann zeta function $\zeta(s)$ and for the character Riemann zeta function $L(\chi, x)$ with real primitive characters satisfying certain positive conditions.

In following discussion, we try to follow the notations used in [1]. The other popular references on special functions is [8]. If we don't mention where a formula comes from, you may find it in the above mentioned references.

2. Preliminaries

Recall that the Euler's $\Gamma(z)$ is defined as

$$\frac{1}{\Gamma(z)} := z\prod_{k=1}^{\infty}\left(1+\frac{z}{k}\right)\left(1+\frac{1}{k}\right)^{-z}, \quad z\in\mathbb{C}, \tag{2.1}$$

and the shifted factorial of $a\in\mathbb{C}$ is defined as

$$(a)_k := \frac{\Gamma(a+k)}{\Gamma(a)}, \quad k\in\mathbb{Z}. \tag{2.2}$$

The Euler's $\Gamma(z)$ satisfies

$$\lim_{n\to\infty}\frac{\Gamma(n+a)}{\Gamma(n+b)}n^{b-a} = 1. \tag{2.3}$$

2000 *Mathematics Subject Classification.* Primary 30E15; Secondary 33D45.

Key words and phrases. Fourier cosine transform; Entire functions; Riemann zeta function; Character Riemann zeta function; Riemann Hypothesis; Ultraspherical polynomials.

The $\Gamma(z)$ function also satisfies the Legendre's duplication formula

$$\Gamma(2z)\Gamma\left(\frac{1}{2}\right) = 2^{2z-1}\Gamma(z)\Gamma\left(z+\frac{1}{2}\right). \tag{2.4}$$

A very useful special value for the $\Gamma(z)$ function is

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}. \tag{2.5}$$

For $\alpha, \beta > -1$ and $n \in \mathbb{Z}^+$, the Jacobi orthogonal polynomials $\left\{P_n^{(\alpha,\beta)}(x)\right\}_{n=0}^{\infty}$ are defined as

$$P_n^{(\alpha,\beta)}(x) := \frac{(\alpha+1)_n}{n!}\sum_{k=0}^{n}\frac{(-n)_k(n+\alpha+\beta+1)_k}{(\alpha+1)_k k!}\left(\frac{1-x}{2}\right)^k. \tag{2.6}$$

The ultraspherical polynomials $\left\{C_n^{(\lambda)}(x)\right\}_{n=0}^{\infty}$ are special cases of the Jacobi polynomials,

$$C_n^{(\lambda)}(x) : = \frac{(2\lambda)_n}{(\lambda+\frac{1}{2})_n}P_n^{(\lambda-\frac{1}{2},\lambda-\frac{1}{2})}(x), \tag{2.7}$$

or

$$C_n^{(\lambda)}(x) = \frac{(2\lambda)_n}{n!}\sum_{k=0}^{n}\frac{(-n)_k(n+2\lambda)_k}{(\lambda+\frac{1}{2})_k k!}\left(\frac{1-x}{2}\right)^k. \tag{2.8}$$

Clearly,

$$C_n^{(\lambda)}(1) = \frac{(2\lambda)_n}{n!}, \tag{2.9}$$

and

$$\frac{C_n^{(\lambda)}(x)}{C_n^{(\lambda)}(1)} = \sum_{k=0}^{n}\frac{n!}{(n-k)!k!}\frac{(n+2\lambda)_k}{(\lambda+\frac{1}{2})_k}\left(\frac{x-1}{2}\right)^k. \tag{2.10}$$

Consequently,

$$c_n^{(\lambda)}(x) := \frac{C_n^{(\lambda)}(x)}{C_n^{(\lambda)}(1)} \geq 1 \tag{2.11}$$

for $x \geq 1$ and $\lambda > 0$. For any $c > 0$ and $\lambda > 0$, the ultraspherical polynomials have the following bound:

$$C_n^{(\lambda)}(\cos(\theta)) = \begin{cases} \theta^{-\lambda}\mathcal{O}(n^{\lambda-1}) & cn^{-1} \leq \theta \leq \frac{\pi}{2} \\ \mathcal{O}(n^{2\lambda-1}) & 0 \leq \theta \leq cn^{-1} \end{cases}, \tag{2.12}$$

which implies

$$c_n^{(\lambda)}(\cos(\theta)) = \frac{C_n^{(\lambda)}(\cos(\theta))}{C_n^{(\lambda)}(1)} = \begin{cases} \theta^{-\lambda}\mathcal{O}(n^{-\lambda}) & cn^{-1} \leq \theta \leq \frac{\pi}{2} \\ \mathcal{O}(1) & 0 \leq \theta \leq cn^{-1} \end{cases}. \tag{2.13}$$

On page 59 of [8] we find the following formula for the Jacobi polynomials,

$$P_{2n}^{(\beta,\beta)}(x) = \frac{(-1)^n n!(\beta+1)_{2n}}{(2n)!(\beta+1)_n}P_n^{(-1/2,\beta)}(1-2x^2), \tag{2.14}$$

which is the same as

$$C_{2n}^{(\beta+1/2)}(x)=\frac{(-1)^n n!}{(\beta+1)_n}\frac{(2\beta+1)_{2n}}{(2n)!}P_n^{(-1/2,\beta)}(1-2x^2). \tag{2.15}$$

The Bessel functions $J_\alpha(z)$ and $I_\alpha(z)$ are defined as

$$J_\alpha(z):=\sum_{k=0}^{\infty}\frac{(-1)^k}{\Gamma(\alpha+k+1)k!}\left(\frac{z}{2}\right)^{2k+\alpha}, \tag{2.16}$$

and

$$I_\alpha(z):=\sum_{k=0}^{\infty}\frac{1}{\Gamma(\alpha+k+1)k!}\left(\frac{z}{2}\right)^{2k+\alpha}, \tag{2.17}$$

respectively, a special case of $J_\nu(z)$ is

$$\cos(z)=\sqrt{\frac{\pi z}{2}}J_{-1/2}(z). \tag{2.18}$$

and exercise 9 on page 235 of [1] gives the inequality

$$|I_\alpha(z)|\le\frac{e^{|z|}}{\Gamma(\alpha+1)}\left(\frac{|z|}{2}\right)^{\alpha},\quad \alpha>-1,\quad z\in\mathbb{C}. \tag{2.19}$$

The Riemann zeta function $\zeta(s)$ is defined as [1, 3, 4, 9]

$$\zeta(s):=\sum_{n=1}^{\infty}\frac{1}{n^s},\quad \Re(s)>1. \tag{2.20}$$

The functions

$$\xi(z):=\frac{z(z-1)}{2\pi^{z/2}}\Gamma\left(\frac{z}{2}\right)\zeta(z), \tag{2.21}$$

and

$$\Xi(z):=\xi\left(\frac{1}{2}+iz\right) \tag{2.22}$$

are entire functions of order 1. They satisfy the following functional equations,

$$\xi(z)=\xi(1-z), \tag{2.23}$$

and

$$\Xi(z)=\Xi(-z). \tag{2.24}$$

The function $\Xi(z)$ has an integral representation

$$\Xi(z)=\int_0^\infty\Phi(t)\cos(zt)dt, \tag{2.25}$$

where

$$\Phi(t):=4\pi\sum_{n=1}^{\infty}\left\{2\pi n^4e^{9t/2}-3n^2e^{5t/2}\right\}\exp\left(-n^2\pi e^{2t}\right). \tag{2.26}$$

It is well known that $\Phi(t)$ is positive, even and fast decreasing on $\mathbb{R}$ [7, 9]. The celebrated Riemann Hypothesis for $\zeta(s)$ is equivalent to the statement that $\Xi(z)$ has only real zeros. More generally, let $\chi(n)$ be a real primitive character to modulus m. The function $L(s,\chi)$ is defined as [1, 3, 4, 6, 9]

$$L(s,\chi):=\sum_{n=1}^{\infty}\frac{\chi(n)}{n^s},\quad \Re(s)>1. \tag{2.27}$$

Let

$$a:=\begin{cases}0 & \chi(-1)=1\\ 1 & \chi(-1)=-1\end{cases}, \tag{2.28}$$

it is known that for

$$\xi(s,\chi|a):=\left(\frac{\pi}{m}\right)^{-(s+a)/2}\Gamma\left(\frac{s+a}{2}\right)L(s,\chi), \tag{2.29}$$

we have

$$\xi(1-s,\chi|a)=\xi(s,\chi|a). \tag{2.30}$$

It is also known that the entire function

$$\Xi(z,\chi|a):=\xi\left(\frac{1}{2}+iz,\chi|a\right) \tag{2.31}$$

is of order 1 and even, and it has an integral representation

$$\Xi(z,\chi|a)=\frac{1}{2}\int_{-\infty}^{\infty}e^{-izt}\Phi(t,\chi|a)dt, \tag{2.32}$$

where

$$\Phi(t,\chi|0):=2e^{t/2}\sum_{n=-\infty}^{\infty}\chi(n)e^{-n^2\pi e^{2t}/m}, \tag{2.33}$$

and

$$\Phi(t,\chi|1):=2e^{3t/2}\sum_{n=-\infty}^{\infty}n\chi(n)e^{-n^2\pi e^{2t}/m}. \tag{2.34}$$

From

$$\sum_{n=-\infty}^{\infty}\chi(n)e^{-n^2\pi x/m}=\frac{1}{\sqrt{x}}\sum_{n=-\infty}^{\infty}\chi(n)e^{-n^2\pi/(mx)},\quad x>0, \tag{2.35}$$

and

$$\sum_{n=-\infty}^{\infty}n\chi(n)e^{-n^2\pi x/m}=x^{-3/2}\sum_{n=-\infty}^{\infty}n\chi(n)e^{-n^2\pi/(mx)},\quad x>0, \tag{2.36}$$

it is easy to verify that

$$\Phi(-t,\chi|a)=\Phi(t,\chi|a),\quad t\in\mathbb{R}. \tag{2.37}$$

The Riemann Hypothesis for $L(s,\chi)$ is equivalent to the statement that $\Xi(z,\chi|a)$ has only real zeros.

3. Main Results

The Bessel function $J_\alpha(z)$ is the scaled large n limit of the Jacobi polynomials [1, 8],

$$\lim_{n\to\infty} n^{-\alpha} P_n^{(\alpha,\beta)}\left(1-\frac{z^2}{2n^2}\right) = \left(\frac{z}{2}\right)^{-\alpha} J_\alpha(z), \tag{3.1}$$

in particular,

$$\lim_{n\to\infty} (n\pi)^{1/2}\, P_n^{(-1/2,\beta)}\left(1-\frac{z^2}{2n^2}\right) = \cos(z), \tag{3.2}$$

and the limits above are uniform on any compact subset of $z \in \mathbb{C}$, as a matter of fact, [1] gives the estimate

$$\left|\frac{n^{-\alpha}}{k!(n-k)!}\frac{(n+\alpha+\beta+1)_k(\alpha+1)_n}{2^{2k}n^{2k}(\alpha+1)_k}\right| \le \frac{C}{k!\Gamma(\alpha+k+1)}, \tag{3.3}$$

for all $0 \le k \le n$ and $n \ge \frac{\alpha+\beta}{2}$, where C is a constant depends on α, β only. Then (3.1) is obtained by applying the Lebesgue dominated convergence theorem. Furthermore, (3.3) also gives the following inequality,

$$\left|n^{-\alpha} P_n^{(\alpha,\beta)}\left(1-\frac{z^2}{2n^2}\right)\right| \le C\sum_{k=0}^{\infty}\frac{|z|^{2k}}{k!\Gamma(\alpha+k+1)} = C\,|z|^{-\alpha}\, I_\alpha\left(2|z|\right) \tag{3.4}$$

for $n \ge \frac{\alpha+\beta}{2}$, where C is a constant depends on α, β only. In the following discussion, we restrict β in the open interval $\beta \in (-\frac{1}{2}, \frac{1}{2})$.

Lemma 3.1. *For $\beta \in (-\frac{1}{2}, \frac{1}{2})$ and sufficiently large $n \in \mathbb{Z}^+$ there is a constant D depends only on β such that*

$$\left|n^{\frac{1}{2}-\beta} C_{4n}^{(\beta+1/2)}\left(\frac{z}{4n}\right)\right| \le D\cosh(2|z|), \tag{3.5}$$

and the

$$\lim_{n\to\infty} (2n)^{1/2-\beta}\,\Gamma\left(\beta+\frac{1}{2}\right) C_{4n}^{(\beta+1/2)}\left(\frac{z}{4n}\right) = \cos(z) \tag{3.6}$$

converges uniformly on any compact subset of $z \in \mathbb{C}$.

Proof. It is a simple calculation with (2.3), (2.4) and (2.5). ☐

The following theorem is an old result of Jensen [5, 7] :

Theorem 3.2. *(Jensen) Let $\Psi(t)$ be a nonnegative even function such that*

1. *$\Psi(t)$ is not identically zero for $t \in \mathbb{R}$,*
2. *$\Psi(t)$ is infinitely differentiable with respect to $t \in \mathbb{R}$,*
3. *For each $n \in \mathbb{Z}^+$,*

$$\lim_{t\to\infty}\frac{\log\left|\Psi^{(n)}(t)\right|}{t} = -\infty. \tag{3.7}$$

4. *The entire function*

$$F(z) = 2\int_0^\infty \Psi(t)\cos(zt)dt \tag{3.8}$$

is of genus at most 1.

Then, the validity of the all following inequalities [5, 7]

$$\int_{-\infty}^{\infty}\int_{-\infty}^{\infty}\Psi(s)\Psi(t)e^{i(s+t)x}(s-t)^{2n}dsdt \geq 0, \quad n\in\mathbb{Z}^{+}, \quad x\in\mathbb{R} \tag{3.9}$$

is necessary and sufficient for $F(z)$ *to have only real zeros.*

By Jensen's necessary and sufficient conditions for $F(z)$ has only real zeros, we only need to verify the inequalities (2.3) are true for all $n\in\mathbb{Z}^{+}$ and all $x\in\mathbb{R}$. Let

$$f_n(x) := \int_{-\infty}^{\infty}\int_{-\infty}^{\infty}\Psi(s)\Psi(t)e^{i(s+t)x}(s-t)^{2n}dsdt \geq 0. \tag{3.10}$$

Since

$$f_n(-x) = f_n(x), \quad x\in\mathbb{R}, \quad n\in\mathbb{Z}^{+}. \tag{3.11}$$

Hence,

$$\begin{aligned} f_n(x) &= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty}\Psi(s)\Psi(t)e^{i(s+t)x}(s-t)^{2n}dsdt \\ &= \int_{-\infty}^{\infty}\int_{-\infty}^{\infty}\Psi(s)\Psi(t)\cos\left((s+t)x\right)(s-t)^{2n}dsdt \\ &= \frac{1}{2}\int_{-\infty}^{\infty}\cos(ux)\left\{\int_{-\infty}^{\infty}\Psi\left(\frac{u+v}{2}\right)\Psi\left(\frac{u-v}{2}\right)v^{2n}dv\right\}du \\ &= \int_{-\infty}^{\infty}\cos(ux)\left\{\int_{0}^{\infty}\Psi\left(\frac{u+v}{2}\right)\Psi\left(\frac{u-v}{2}\right)v^{2n}dv\right\}du \\ &= 2\int_{0}^{\infty}\cos(ux)\left\{\int_{0}^{\infty}\Psi\left(\frac{u+v}{2}\right)\Psi\left(\frac{u-v}{2}\right)v^{2n}dv\right\}du, \end{aligned} \tag{3.12}$$

where we have made the simple change of variables

$$u = s+t, \quad v = s-t, \tag{3.13}$$

and used the fact $\Psi(t)$ is an even function.

Theorem 3.3. *Let* $\Psi(t)$ *and* $F(z)$ *be as defined as in Theorem 3.2 and satisfy all the four conditions of Theorem 3.2 plus the condition that* $\Psi(t)$ *is positive almost everywhere on* $\mathbb{R}$*. Then* $F(z)$ *has only real zeros.*

Proof. From the Jensen's result, 3.2, it is enough for us to prove (3.9). From (3.12) we only need to show that

$$\int_{0}^{\infty}\cos(ux)\left\{\int_{0}^{\infty}\Psi\left(\frac{u+v}{2}\right)\Psi\left(\frac{u-v}{2}\right)v^{2n}dv\right\}du \geq 0, \quad n\in\mathbb{N}, \quad x>0. \tag{3.14}$$

Lemma 3.1 and condition 2, 3 of Theorem 3.2 and an application of the Lebesgue's Dominated Convergence Theorem show that (3.15) is same as for some $\beta\in(-\frac{1}{2},\frac{1}{2})$,

$$\lim_{N\to\infty}\frac{\Gamma\left(\beta+\frac{1}{2}\right)}{(2N)^{\beta-\frac{1}{2}}}\int_{0}^{\infty}C_{4N}^{(\beta+\frac{1}{2})}\left(\frac{ux}{4N}\right)\left\{\int_{0}^{\infty}\Psi\left(\frac{u+v}{2}\right)\Psi\left(\frac{u-v}{2}\right)v^{2n}dv\right\}du \geq 0 \tag{3.15}$$

for all $n \in \mathbb{N}$ and $x > 0$. Since the limit (3.15) exists, then it is sufficient to show that for some $\beta \in (-\frac{1}{2}, \frac{1}{2})$, there infinitely many large $N \in \mathbb{N}$ such that

$$\int_0^\infty C_{4N}^{(\beta+\frac{1}{2})}(ux) \left\{ \int_0^\infty \Psi\left(\frac{u+v}{2}\right) \Psi\left(\frac{u-v}{2}\right) v^{2n} dv \right\} du \geq 0 \tag{3.16}$$

for all fixed $x > 0$ and all fixed $n \in \mathbb{N}$. Consider

$$g_N(x) := \int_0^\infty c_{4N}^{(\beta+\frac{1}{2})}(ux) \left\{ \int_0^\infty \Psi\left(\frac{u+v}{2}\right) \Psi\left(\frac{u-v}{2}\right) v^{2n} dv \right\} du, \tag{3.17}$$

then

$$\begin{aligned} g_N(x) &= \int_0^{\frac{1}{x}} c_{4N}^{(\beta+\frac{1}{2})}(ux) \left\{ \int_0^\infty \Psi\left(\frac{u+v}{2}\right) \Psi\left(\frac{u-v}{2}\right) v^{2n} dv \right\} du \\ &\quad + \int_{\frac{1}{x}}^\infty c_{4N}^{(\beta+\frac{1}{2})}(ux) \left\{ \int_0^\infty \Psi\left(\frac{u+v}{2}\right) \Psi\left(\frac{u-v}{2}\right) v^{2n} dv \right\} du \\ &= I_1 + I_2. \end{aligned} \tag{3.18}$$

Formula (2.11) implies that

$$I_2 \geq \int_{\frac{1}{x}}^\infty \int_0^\infty \Psi\left(\frac{u+v}{2}\right) \Psi\left(\frac{u-v}{2}\right) v^{2n} dv du > 0 \tag{3.19}$$

for all $x > 0$ and all $n \in \mathbb{N}$. Make the change of variables

$$ux = \cos(\theta), \quad 0 < \theta < \frac{\pi}{2} \tag{3.20}$$

in I_1 to get

$$\begin{aligned} I_1 = x^{-1} &\int_0^{\frac{\pi}{2}} c_{4N}^{(\beta+\frac{1}{2})}(\cos(\theta)) \sin(\theta) d\theta \\ &\times \left\{ \int_0^\infty \Psi\left(\frac{x^{-1}\cos(\theta)+v}{2}\right) \Psi\left(\frac{x^{-1}\cos(\theta)-v}{2}\right) v^{2n} dv \right\}. \end{aligned} \tag{3.21}$$

Clearly, by (2.13) we have

$$\begin{aligned} |I_1| &\leq \frac{a(x)}{x} \int_0^{\frac{\pi}{2}} \left| c_{4N}^{(\beta+\frac{1}{2})}(\cos(\theta)) \sin(\theta) \right| d\theta \\ &= \mathcal{O}\left(N^{-\beta-\frac{1}{2}}\right), \end{aligned} \tag{3.22}$$

where

$$a(x) := \sup_{0 \leq \theta \leq \pi/2} \left\{ \int_0^\infty \Psi\left(\frac{x^{-1}\cos(\theta)+v}{2}\right) \Psi\left(\frac{x^{-1}\cos(\theta)-v}{2}\right) v^{2n} dv \right\} < \infty. \tag{3.23}$$

Hence, there is a positive integer N_0, for each $N > N_0$ we have

$$|I_1| \leq \frac{1}{2} \int_{\frac{1}{x}}^\infty \int_0^\infty \Psi\left(\frac{u+v}{2}\right) \Psi\left(\frac{u-v}{2}\right) v^{2n} dv du, \tag{3.24}$$

consequently, for each $N > N_0$ we have

$$g_N(x) \geq I_2 - |I_1| \geq \frac{1}{2} \int_{\frac{1}{x}}^\infty \int_0^\infty \Psi\left(\frac{u+v}{2}\right) \Psi\left(\frac{u-v}{2}\right) v^{2n} dv du > 0, \tag{3.25}$$

and the conclusion of our Theorem 3.3 follows. □

Theorem 3.3 give the validation of the Riemann Hypothesis:

Corollary 3.4. *The Riemann Hypothesis for $\zeta(s)$ is true. The Riemann Hypothesis for $L(s,\chi)$ is true for real primitive characters χ satisfying $\Phi(t,\chi|a)\geq 0$.*

As it is remarked in [2], one could prove that $\Phi(t,\chi|a)\geq 0$ when the modulus is small, and the condition may be false for certain modulus. One may speculate that the corresponding Riemann Hypothesis are false for some of those $L(s,\chi)$ which fails the positive condition.

Acknowledgement. This work is partially supported by Chinese National Natural Science Foundation grant No.10761002, Guangxi Natural Science Foundation grant No.0728090.

References

[1] G. E. Andrews, R. Askey and R. Roy, *Special Functions,* Cambridge University Press, Cambridge, 1999.

[2] Philippe Biane, Jim Pitman and Marc Yor, Probability Laws Related to the Jacobi Theta and Riemann Zeta Functions, and Brownian Excursions, Bulletin (New Series) of the American Mathematical Society, Volume 38, Number 4, Pages 435-465.

[3] H. Davenport, Multiplicative Number Theory, Springer-Verlag, New York,

[4] A. Erdélyi, *Higher Transcendental Functions,* Vol.I, Vol.II, Vol.III, Robert E. Krieger Publishing Company, Malabar, Florida, 1985.

[5] George Gasper, Using sums of squares to prove that certain entire functions have only real zeros, dedicated to the memory of Ralph P. Boas, Jr. (1912–1992), published in *Fourier Analysis: Analytic and Geometric Aspects*, W.O. Bray, P.S. Milojevic and C.V. Stanojevic, eds., Marcel Dekker, 1994, pp. 171–186.

[6] L. K. Hua, *Introduction to Number Theory*, Springer-Verlag, New York, 1982.

[7] G. Pólya, Über die algebraisch-funktionentheoretischen Untersuchungen von J. L. W. V. Jensen, *Kgl. Danske Videnskabernes Selskab. Math.-Fys. Medd.* 7 (17) (1927), pp. 3–33; reprinted in his *Collected Papers*, Vol. II, pp. 278–308.

[8] G. Szegö, *Orthogonal Polynomials*, AMS Colloquium Publications, Vol.23, 4th. edition, Providence, Rhode Island, 1975.

[9] E. C. Titchmarsh, *The Theory of Riemann Zeta Function*, second edition, Clarendon Press, New York, 1987.

[10] G. N. Watson, *A Treatise on the Theory of Bessel Functions*, Cambridge University Press, London, 1922.

Current address: School of Mathematical Sciences, Guangxi Normal University, Guilin City, Guangxi 541004, P. R. China.

E-mail address: `ruimingzhang@yahoo.com`